
· · · · · · · · · · · · · · · · · · ·		
.5', 1., 1 1.,		
· · · · ·		
$\frac{h}{1} = \frac{1}{1} \frac{q_{1}}{1} + \frac{q_{1}}{1$		
ν ^p , <u>γ</u> , ρ _λ , γ, γ, γ, α ι γ,		
a h c h f f f h h h		
h h h a h h a h		
a the second sec		
fa af a graph		
a li i a ja a i i i a		
• _ • _ •		
$-r^{\alpha}$ η η η η η η η		
fa h a fa h		
, The area of the		
$i = 2T^{-1} (1 A^{-1} (1A^{-1} (1A$		
alia para a th		
ρ _{, γ} -, -, α, γ,		
	$ \begin{array}{c} \underline{\mathbf{h}} & \mathbf{zq} \\ \mathbf{q}_{1,1,1,1} \mathbf{z}_{1}^{\mathbf{p}} & \underline{\mathbf{x}}_{1,1} & 10, \\ \mathbf{h} & \mathbf{z}_{1,1,1} \mathbf{z}_{1,1} & \mathbf{z}_{1,1} & \mathbf{z}_{1,1} \\ \mathbf{z}_{1,1,1} & \mathbf{z}_{1,1} & \mathbf{z}_{1,1} & \mathbf{z}_{1,1} \\ \mathbf{z}_{1,1,1} \\ \mathbf{z}_{1,1,1} & \mathbf{z}_{1,1} \\ \mathbf{z}_{1,1,1} \\ \mathbf{z}_{1$	
	h at a tra tra tra h	
	$\frac{10}{10}$	$\frac{12}{2} - \frac{12}{2} $
	z 10. (,).	$-\frac{1}{2q} - 2q = 2,$
	z 10. (). z d q q z b d z	
μ , ε α <u></u> ε	$\begin{array}{c} q \\ p \\$	$\begin{array}{c} \mathbf{a} & \mathbf{a} & \mathbf{r} \\ $
$\frac{\mathbf{z}}{\mathbf{z}} \stackrel{\mathbf{h}}{\longrightarrow} \begin{array}{c} \mathbf{c} \mathbf{c} \mathbf{c} \mathbf{c} \mathbf{c} \mathbf{c} \mathbf{c} \mathbf$	$a \rightarrow a + b \rightarrow $	$\frac{1}{(2\pi)^2} + \frac{1}{2} +$
	a a har har a dry har a	$\begin{array}{c} \begin{array}{c} g & a \\ g & a \\ \hline \end{array} \\ \begin{array}{c} g & a \\ \end{array} \\ \begin{array}{c} g & a \\ \end{array} \\ \begin{array}{c} g & a \\ \end{array} \end{array} \\ \begin{array}{c} g & a \\ \end{array} \end{array} \end{array} \\ \begin{array}{c} g & a \\ \end{array} \end{array} \end{array} $ \\ \begin{array}{c} g & a \\ \end{array} \end{array} \end{array} \\ \begin{array}{c} g & a \\ \end{array} \end{array} \end{array} \\ \begin{array}{c} g & a \\ \end{array} \end{array} \end{array} \\ \begin{array}{c} g & a \\ \end{array} \end{array} \end{array} \\ \begin{array}{c} g & a \\ \end{array} \end{array} \end{array} \\ \end{array} \end{array} \\ \begin{array}{c} g & a \\ \end{array} \end{array} \end{array} \\ \begin{array}{c} g & a \\ \end{array} \end{array} \end{array} \end{array} \\ \end{array} \end{array} \\ \begin{array}{c} g & a \\ \end{array} \end{array} \end{array} \end{array} \\ \end{array} \end{array} \end{array} \\ \end{array} \end{array} \\ \begin{array}{c} g & a \\ \end{array} \end{array} \end{array} \end{array} \\ \end{array} \end{array} \end{array} \\ \begin{array}{c} g & a \\ \end{array} \end{array} \end{array} \\ \end{array} \end{array} \\ \end{array} \end{array} \end{array} \\ \end{array} \end{array} \\ \end{array} \end{array} \end{array} \end{array}
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		a a b f f
<u>, pr pr. , pr.</u>	- I Correction of the second	$\mathbf{z} \mathbf{q} = \mathbf{z} \mathbf{q} + \mathbf{z} \mathbf{z} + \mathbf{z} + $
	a a a b b b b a b a b a b a b a b a b a	$\begin{array}{cccccccccccccccccccccccccccccccccccc$
• • • • • • • • • • • • • • • • • • •		it to construct the second


$\begin{array}{c} 2 \\ p \\ p \\ c \\ a \\ b \\ c \\ c$
 01, , , , , , , , , , , , , , , , , , ,
$\begin{array}{c} 2 & b \\ r & f \\ r & f \\ q \\ r \\ r$



-

to a star i partition

1-



	<u> 1, </u>	
	<u>2 - 79 12</u> 9 7~ 79	
	, <u>a</u> a <u>b</u> <u>c</u>	
	$A_{1,1}^{p}$, A_{2}^{p} , A_{1}^{p} , A_{1}^{p} , A_{2}^{p} , A	
	f for F mrs , g b	
	سر) a طریکور عن تعد مطرسر عن و	
	J raling all mrs par a	
	halder arrent g	
	Afra a for a ref in	
	All the the second second	
	$\mathbf{p} = \left\{ \begin{array}{ccc} \mathbf{p} & \mathbf{p} \\ \mathbf{p} \\ \mathbf{p} & \mathbf{p} \\ \mathbf{p} \\ \mathbf{p} & \mathbf{p} \\ \mathbf{p} & \mathbf{p} \\ \mathbf{p} & \mathbf{p}$	
	hallon, a lith al ch	
···		
h h	a a p a p a p	, , ,
	$\begin{array}{c} 2 \\ a \\ a \\ a \\ a \\ a \\ b \\ b \\ a \\ b \\ c \\ c$	$p = \frac{p}{1 + \frac{p}{1$
a set of a set of the set and set		ita in ha a in hating a free
N , F , F , A , A , A , A , A , A , A , A		
	at a as stands when the set of the	a a h a b
$1, \dots - 2q 2.$	ar a f i c f a i c h	acail acail har
$\mathbf{A}_{\mathbf{A}} = \mathbf{A}_{\mathbf{A}} \mathbf{A} \mathbf{A}} \mathbf{A}_{\mathbf{A}} A$		م ي م الم <u>الم م م م م م م م م م م م</u>
		r a da krak, r a k
∠Q ,		$\begin{array}{c} & & & & & & & & & & & & & \\ & & & & & $
⁰ , z [*] ₂ , zz [*] q _{*i} s _i , , , , (, ")- ₂q , 2.		$f_{1} \qquad , \qquad a_{1} \in \mathcal{C} a_{1} \qquad f_{2} b_{1} = \frac{b_{1}}{2}$

1 Q, ha la h h aa g 🖉 pr v p × . a ar; ra, ar; c, b a ar; ra, ar; c, b p a ar; ra, ar; c, b a la b b b b a b The second secon i me e e e fa ha man ha i a me ha man ha i a me ha ha i a me ha i a he i a ha i a he i a a 1 f q , c , c , b f , b , c , b , c , b a , q , c , c , b 10. ()()(-,), b , -, q , b a a b a b a ą, 3 , 1, βq, q_ρ⁰, μ_μ², μ 00 17- 1 P 1 1- 11, q mo to the total to the - 17 S $\begin{array}{c} \mathbf{q} \\ \mathbf{p} \\ \mathbf{h} \\ \mathbf{h} \\ \mathbf{h} \\ \mathbf{h} \\ \mathbf{q} \\ \mathbf{r} \\ \mathbf{h} \\ \mathbf{h} \\ \mathbf{h} \\ \mathbf{q} \\ \mathbf{r} \\ \mathbf{h} \\ \mathbf{h} \\ \mathbf{h} \\ \mathbf{q} \\ \mathbf{r} \\ \mathbf{h} \\ \mathbf{$ þ . 10, Ça, a,Ç , 17, , 00 . 5 20 0. b ું. કો , ۶, $\begin{array}{c} \mathbf{q} \\ \mathbf{$ f. ar 1 $\begin{array}{c} \mathbf{q} \\ \mathbf{$ 6 * 72<mark>0</mark> p a wap f, $([a], [b]_{W_{c}}, [b]_{W_{$ <u>م</u>ر ب -a h · · · · · · · · · · · · ·

þ

∎ a,c

1, 27

1 20. (,^h)(:

Q 129 .

\$

ba a cart

,́άþ, ,⊈,))¢,

· · · · ·

Q

⁻Q h f